

Agricultural Marketing Service, USDA

§ 29.1163

Unripe, tight leaf structure, heavy, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

B3KK Good Quality Excessively Scorched Leaf

Unripe, close leaf structure, heavy, normal width, 16 inches (40.6 cm) or over in length. Uniformity, 80 percent; injury tolerance, 15 percent.

B4KK Fair Quality Excessively Scorched Leaf

Unripe, close leaf structure, heavy, normal width. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

B5KK Low Quality Excessively Scorched Leaf

Unripe, tight leaf structure, heavy, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

B6KK Poor Quality Excessively Scorched Leaf

Unripe, tight leaf structure, heavy, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

B4KV—Fair Quality Variegated Greenish Leaf

Unripe, firm leaf structure, medium body, normal width. Uniformity, 70 percent; tolerance, 25 percent waste.

B5KV—Low Quality Variegated Greenish Leaf

Unripe, firm leaf structure, medium body, narrow. Uniformity, 70 percent; tolerance, 30 percent waste.

B6KV—Poor Quality Variegated Greenish Leaf

Unripe, firm leaf structure, medium body, stringy. Uniformity, 70 percent; tolerance, 40 percent waste.

B3S—Good Quality Slick Leaf

Unripe, close leaf structure, fleshy, normal width, 16 inches or over in length. Uniformity, 80 percent; injury tolerance, 15 percent.

B4S—Fair Quality Slick Leaf

Unripe, close leaf structure, fleshy, normal width. Uniformity, 70 percent; injury tolerance, 20 percent, of which not over 5 percent may be waste.

B5S—Low Quality Slick Leaf

Unripe, tight leaf structure, fleshy, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

B4G—Fair Quality Green Leaf

Immature, close leaf structure, fleshy, oily, normal width. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

B5G—Low Quality Green Leaf

Immature, tight leaf structure, fleshy, lean in oil, narrow. Uniformity, 70 percent; injury

tolerance 30 percent, of which not over 10 percent may be waste.

B6G—Poor Quality Green Leaf

Immature, tight leaf structure, fleshy, lean in oil, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

B5GR—Low Quality Green Red Leaf

Immature, tight leaf structure, heavy, lean in oil, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

B4GK—Fair Quality Green Variegated Leaf

Immature, close leaf structure, heavy, normal width. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

B5GK—Low Quality Green Variegated Leaf

Immature, tight leaf structure, heavy, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

B6GK—Poor Quality Green Variegated Leaf

Immature, tight leaf structure, heavy, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

B5GG—Low Quality Gray Green Leaf

Immature, tight leaf structure, heavy, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

[42 FR 21092, Apr. 25, 1977, as amended at 48 FR 29671, June 28, 1983; 49 FR 16756, Apr. 20, 1984; 51 FR 25028, July 10, 1986]

§ 29.1163 Smoking Leaf (H Group).

This group consists of leaves normally grown at or above the midportion of the stalk. Leaves of the H group show a high degree of maturity, more open leaf structure in relation to the B Group, and a material amount of injury characteristic of very ripe leaf tobacco.

Grades, Grade Names, Minimum Specifications, and Tolerances

H3F—Good Quality Orange Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, strong color intensity, normal width, 16 inches or over in length. Uniformity, 80 percent; injury tolerance, 15 percent.

H4F—Fair Quality Orange Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, moderate color intensity, normal width. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

H5F—Low Quality Orange Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, weak color intensity, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

§ 29.1164

7 CFR Ch. I (1-1-03 Edition)

H6F—Poor Quality Orange Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, weak color intensity, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

H4FR—Fair Quality Orange Red Smoking Leaf

Mellow, open leaf structure, fleshy, lean in oil, moderate color intensity, normal width. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

H5FR—Low Quality Orange Red Smoking Leaf

Mellow, open leaf structure, fleshy, lean in oil, weak color intensity, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

H6FR—Poor Quality Orange Red Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, weak color intensity, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

H4K—Fair Quality Variegated Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, moderate color intensity, normal width. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

H5K—Low Quality Variegated Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, weak color intensity, narrow. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

H6K—Poor Quality Variegated Smoking Leaf

Mellow, open leaf structure, medium body, lean in oil, weak color intensity, stringy. Uniformity, 70 percent; injury tolerance 40 percent, of which not over 20 percent may be waste.

[42 FR 21092, Apr. 25, 1977, as amended at 48 FR 29671, June 28, 1983; 49 FR 16756, Apr. 20, 1984; 51 FR 25028, July 10, 1986]

§ 29.1164 Cutters (C Group).

This group consists of leaves normally grown at or just below the midportion of the stalk. Leaves of the C group have a tendency to roll concealing the stem or midrib. Cutters usually have a rounded tip, are thin to medium in body, and show some ground injury.

Grades, Grade Names, Minimum Specifications, and Tolerances

C1L—Choice Quality Lemon Cutters

Ripe, open leaf structure, medium body, oily, deep color intensity, spready, 20 inches or over in length. Uniformity, 90 percent, injury tolerance, 5 percent.

C2L—Fine Quality Lemon Cutters

Ripe, open leaf structure, thin, oily, deep color intensity, spready, 20 inches or over in length. Uniformity, 85 percent; injury tolerance, 10 percent.

C3L—Good Quality Lemon Cutters

Ripe, open leaf structure, thin, oily, strong color intensity, spready, 18 inches or over in length. Uniformity, 80 percent; injury tolerance, 15 percent.

C4L—Fair Quality Lemon Cutters

Ripe, open leaf structure, thin, lean in oil, moderate color intensity, normal width, 16 inches or over in length. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

C5L—Low Quality Lemon Cutters

Ripe, open leaf structure, thin, lean in oil, weak color intensity, normal width, 16 inches or over in length. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

C4LL—Fair Quality Whitish-Lemon Cutters

Unripe, firm leaf structure, thin (papery), lean in oil, normal width, 16 inches or over in length. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

C5LL Low Quality Whitish-Lemon Cutters

Unripe, firm leaf structure, thin (papery), lean in oil, normal width, 16 inches (40.6 cm) or over in length. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

C5LP—Low Quality Lemon Cutters (Primings Side)

Prematurely ripe, open leaf structure, thin, lean in oil, pale color intensity, normal width, 16 inches or over in length. Uniformity, 70 percent; injury tolerance 30 percent, of which not over 10 percent may be waste.

C1F—Choice Quality Orange Cutters

Ripe, open leaf structure, medium body, oily, deep color intensity, spready, 20 inches or over in length. Uniformity, 90 percent; injury tolerance, 5 percent.

C2F—Fine Quality Orange Cutters

Ripe, open leaf structure, medium body, oily, deep color intensity, spready, 20 inches or over in length. Uniformity, 85 percent; injury tolerance, 10 percent.

C3F—Good Quality Orange Cutters

Ripe, open leaf structure, medium body, oily, strong color intensity, spready, 18 inches or over in length. Uniformity, 80 percent; injury tolerance, 15 percent.

C4F—Fair Quality Orange Cutters

Ripe, open leaf structure, medium body, lean in oil, moderate color intensity, normal width, 16 inches or over in length. Uniformity, 70 percent; injury tolerance 20 percent, of which not over 5 percent may be waste.

C5F—Low Quality Orange Cutters

Ripe, open leaf structure, medium body, lean in oil, weak color intensity, normal